




# COTMAN™ Cotton Management System

COTMAN is divided into 2 components:

1. **SQUAREMAN** is used to monitor crop development up to time of first flowers. It is used to monitor square retention and plant stress.
2. **BOLLMAN** is used when the crop is flowering to monitor boll loading and plant stress and to help with end-of-season crop termination decisions. BOLLMAN utilizes NAWF data.



For a cotton crop to follow the COTMAN target development curve, the crop should begin squaring 35 days after planting, and first flowers should appear about 25 days later, around 60 days after planting. Physiological cutout is expected approximately 80 days after planting when the crop averages NAWF = 5.



# COTMAN™ Data Collection




Use a T-stick to take stand counts.



Make determination of FN when squares consistently can be found across the field.

Once per season you enter these data into the COTMAN program.

1. Farm information - total acres, harvest capacity, weather risk...
2. Field information - field acreage, variety, planting date, row spacing...
3. Stand count – the number of plants in 3 row feet across 24 consecutive rows at 4 locations per field.
4. First fruiting node (FN) - the average first fruiting node (1<sup>st</sup> sympodial branch) on 10 plants at 4 locations in the field.



Count from the cotyledons (node “0”) up to the first sympodial branch to determine FN.




# SQUAREMAN Data Collection

**Weekly Data Collection:** Begin collecting SQUAREMAN data when the crop starts squaring (you may wish to collect FN data at the same time you start collecting SQUAREMAN data). Repeat the following procedure in 4 different locations in each field:


1. Measure the average plant height.
2. From 5 consecutive plants, start at the 1st fully expanded true leaf in the terminal and check for the presence or absence of 1st position squares. Record a “1” if a square is present and enter a “0” if the square has shed.
3. Repeat steps 1 & 2 on the adjacent row.



**Measure plant height from the soil to the terminal in inches – 1 measurement on both rows at each site.**



**Check for presence or absence of 1st position squares starting the top of the plant with the 1st fully expanded leaf.**



**Don't start checking for squares on this node because this leaf is not yet unfurled - its leaf tips are touching.**

**Check for presence or absence of 1<sup>st</sup> position squares starting at the 1st node with an unfurled leaf and continue down the plant.**


Deciding where to sample in the field ....:

- ✓ Choose the 4 sample sites where plant growth is typical for the field.
- ✓ Try to sample the same general areas in the same order every week.
- ✓ If the field is irrigated, always stay inside the irrigated area.
- ✓ It is a good idea to ask the grower where to make your counts in the fields because he will know the fields best..



# SQUAREMAN Mapper Tips


1. Start at the top of the plant at the 1st fully expanded true leaf – its tips should not be touching.
2. Don't damage the terminal area digging around for small squares.
3. If the 1<sup>st</sup> square is dark or obviously damaged record a "0". If the square looks normal record a "1".
4. Check the remaining 1<sup>st</sup> position squares down the plant until you reach a monopodial branch.
5. Move to the next plant and begin again.



## Follow the Stipule Rule:

Stipules are small leaves on each side of a square. It will save time looking for the square if you train your eye to look precisely between the stipules. The stipules are pointing where you need to look.

A pair of stipules on a sympodial branch will have between them either 1) a square or boll or 2) a scar where a square or boll has been shed



For this plant you would record



1 <sup>st</sup> Position Square on Node	Record on Psion Workabout
1	1
2	1
3	0
4	1
5	1
6	0
7	1

1 = square present  
 2 = boll present  
 0 = square or boll shed



# BOLLMAN Data Collection

When first flowers appear, start collecting NAWF data. Repeat the following procedure at 4 locations per field:


1. Count the number of nodes **above** the uppermost 1st position white flower (NAWF) from 5 plants in one row and 5 plants from an adjacent row. When the crop is just starting to flower, you may have to “swim” down the row to find 5 plants with white flowers.
2. When counting, stop at the uppermost unfurled leaf in the terminal (*DO NOT COUNT A LEAF THAT IS NOT YET UNFURLED*).
3. Stop collecting BOLLMAN data when the field averages a NAWF value of 5.




Fructing Positions:  
(From Main-Stem)

Select only plants with a **WHITE** flower on the **FIRST** fructing position. On this branch there is a pink flower on the 2<sup>nd</sup> position so it would not be selected.

**NAWF = Nodes Above White Flower**




The NAWF count for this plant is 5.



Do not count this node - this leaf is not yet unfurled - its leaf tips are touching.  
This leaf is unfurled and so this node would be the last node counted when determining NAWF.

Plant mappers must understand plant structure.

**LEAVES:**



(A) Cotyledon leaves, (B) first true leaf, (C) & (D) true leaves from main-stem and fruiting branch.



Cotyledons are the 1st leaves to emerge, and they are the only leaves side-by-side on the main-stem.





1st true leaf (Heart shaped)  
True leaves (lobed leaves)

Cotyledon - Bean shaped & lies side-by-side on the main-stem.

**Main Stem Branches:**

1. After a sympodial branch (fruiting branch) produces a square it will begin to grow again and will zig-zag.
2. A monopodial branch (vegetative branch) is a continuous limb.
3. Cotyledons lie side-by-side on the main-stem. All other branches rotate up the plant.



Sympodial branch (zig-zag )

Monopodial branches look like individual plants growing from the main-stem, and they can eventually set fruit.